

**MARK SCHEME for the May/June 2011 question paper
for the guidance of teachers**

5054 PHYSICS

5054/21

Paper 2 (Theory), maximum raw mark 75

This mark scheme is published as an aid to teachers and candidates, to indicate the requirements of the examination. It shows the basis on which Examiners were instructed to award marks. It does not indicate the details of the discussions that took place at an Examiners' meeting before marking began, which would have considered the acceptability of alternative answers.

Mark schemes must be read in conjunction with the question papers and the report on the examination.

- Cambridge will not enter into discussions or correspondence in connection with these mark schemes.

Cambridge is publishing the mark schemes for the May/June 2011 question papers for most IGCSE, GCE Advanced Level and Advanced Subsidiary Level syllabuses and some Ordinary Level syllabuses.

Page 2	Mark Scheme: Teachers' version	Syllabus
	GCE O LEVEL – May/June 2011	5054

Do not accept fractions. No penalty for 2 sig. fig. or for 1 sig. fig. where exactly correct.
Only one unit and only one fraction penalty per question.

Section A

- 1 (a) (i) 11.5 m/s B1
- (ii) decrease in speed // negative gradient C1
equal changes/decreases in speed in the same time // const. neg. grad. on v-t graph A1
- (b) (i) flat line at 18 m/s from $t = 0$ to 15 B1
constant slope downwards parallel to initial line (by eye) B1
- (ii) greater area **under** graph // higher **initial/average** speed B1 [6]
- 2 (a) (i) X weight // (force of) gravity // gravitational (force) B1
and Y air resistance // (air) drag // wind resistance // **air** friction B1
- (ii) (Y) opposes motion // diver moves down // air molecules hit faster from below B1
- (b) (i) accelerates // falls faster // speed/velocity greater B1
X is larger than Y // resultant/net force downwards B1
- (ii) **two** of: B2 [6]
X stays the same // Y increases // forces become equal (and opposite)
- 3 (a) (i) 7000 (J) seen **or** 50% **used** somewhere C1
 $(P =) E/t // 14000/t // 7000/t$ seen // 7000 J/minute // 420 000 J/hour C1
120 W A1
- (ii) water after hitting turbine still moves // has KE/energy/velocity // energy lost due to friction // friction **and** location // heat/internal energy **and** location // water misses turbine B1
- (b) (i) can be replaced/made // will not run out B1
- (ii) coal, oil, gas, peat, nuclear, uranium (**not** solar) B1 [6]

www.PapaCambridge.com

Page 3	Mark Scheme: Teachers' version	Syllabus
	GCE O LEVEL – May/June 2011	5054

- 4 (a) $(E/Q =) mc\Delta T$ in any algebraic or numerical form e.g. 4200×16 , 2100×5
 $67\ 200$ or $10\ 500$ or $77\ 700$ seen or $(E/Q =) ml$ or mL algebraic seen
 4.2×10^5 J A1
- (b) (i) break bonds // separate molecules // give molecules more P.E. B1
- (ii) (different) change in distance // molecules not so far apart // incomplete bond
 breaking // doesn't push atmosphere back // less work against atmosphere B1 [5]
- 5 (a) gives out (visible) light // glows B1
 when hit by uv/electrons or spark/discharge/current in tube B1
- (b) X-rays or gamma B1
- (c) $(f =) v/\lambda$ numerical or any algebraic form, e.g. $v = f\lambda$ C1
 8.3×10^{14} Hz A1 [5]
- 6 (a) reflection (of sound/ultrasound) B1
- (b) waves of same period (by eye) B1
 smaller amplitude (by eye) B1
- (c) (i) 20–20 000 Hz B1
- (ii) higher than (i) B1 [5]
- 7 (a) upward arrow (**not** curved) on iron bar B1
- (b) attraction/force not enough//weight of bar too high//friction at pivot//with copper bar B1
- (c) **three** of:
 electromagnet works // magnetic field created
 iron bar moves/lifted up
 spring pulls copper bar across/contracts
 contacts break circuit // contacts open B3
- (d) more turns in coil // more iron in electromagnet // electromagnet nearer iron bar //
 iron bar less weight // weaker spring M1
 correct explanation which involves force on iron bar A1 [7]

Page 4	Mark Scheme: Teachers' version	Syllabus
	GCE O LEVEL – May/June 2011	5054

- 8 (a) all three correct: force, field, current
- (b) (i) Fd // 20×4 ; 20×2 ; 20×0.04 ; 20×0.02 ; $20 \times 4 \times 2$; 160
80 N cm, 0.8 N m C1
A1
- (ii) more turns (on coil)
use soft iron
more current
increase AB or CD
increase BC or AD ANY 2 B1 [5]

Section B

- 9 (a) (i) d.c. current flows in one direction or
a.c. current flows in one direction then the other B1
- (ii) mention of magnetic field/flux B1
(magnetic) field lines // flux cuts coil // flux changes in coil B1
induction of voltage/current B1
something **relevant** reverses (e.g. field/flux cuts in one direction then the other // N pole approaches then leaves // N pole approaches and S pole approaches) B1
and link to a.c. B1
- (iii) **two** of: thicker wires; more turns of coil; stronger magnet; faster rotation; lower resistance (of lamp) B2
- (b) (i) to reduce heat/energy/power loss (on the power lines) B1
(higher voltage means) lower current B1
- (ii) 25:400 // 1:16 // 0.0625 B1
- (iii) reduces resistance // less power loss // costs less to run // more current // more power B1
increases weight // more support needed // more wind resistance // more ice forms // costs more to install B1
- (iv) 13 A B1
($I = P/V$) // 4.2 A // 4.17 etc A B1
must choose higher value to avoid fuse blowing // other fuses melt B1 [15]

Page 5	Mark Scheme: Teachers' version	Syllabus
	GCE O LEVEL – May/June 2011	5054

- 10 (a) (i) brown and green
red
- (ii) 99×10^9 or $9.9 \times 10^{10} \Omega$ B1
- (iii) *less*: (likely to) burn out/blow // become too hot B1
greater: (likely to) be large (in size) B1
- (b) (i) both involve energy and charge // measured in J/C/volts/by voltmeter B1
energy change is from other forms (accept chemical) to electrical in e.m.f. B1
energy change is from electrical to other forms (accept heat/light) in p.d. B1
(or e.m.f. is property of source and p.d. is property of (part of) circuit B2)
- (ii) correct symbol B1
correct direction B1
- (iii) p.d. reduces/(approximately) constant **and** current reduces B1
- (iv) $R = V/I$ in any form, e.g. 1.7/0.025, 1.7/25 C1
correct conversion to mA, e.g. 0.025/7.3 seen C1
68 or 360 seen C1
290 Ω , 292 Ω A1 [15]
- 11 (a) (i) 24 B1
- (ii) charges (of electrons and protons) cancel // protons and electrons have the same (size of) charge but opposite in sign B1
- (b) (i) nuclear at start // nucleus loses energy B1
to kinetic/e.m. energy (condone light/photon/ γ) B1
- (ii) beta 0 at top B1
beta -1 at bottom B1
Mg 24 at top B1
Mg 12 at bottom B1
- (c) (i) 600 at 15 hours B1
300 at 30 hours B1
line of decreasing gradient (**not** if it cuts time axis before $t = 30$) B1
- (ii) emission is random // not predictable // not regular // exptl. error B1
- (iii) 4800 (counts per minute) // 80 counts /sec B1
- (iv) GM tube/solid state detector/cloud chamber M1
ratemeter/data logger // counter/scalar **and** stopwatch/timer A1 [15]