Cambridge Pre-U

CANDIDATE NAME					
CENTRE NUMBER			CANDIDATE NUMBER		

0123456789

BIOLOGY 9790/03

Paper 3 Case Study and Synoptic Essay

For examination from 2020

SPECIMEN PAPER

1 hour 45 minutes

You must answer on the question paper.

No additional materials are needed.

INSTRUCTIONS

- Section A: answer all questions.
- Section B: answer one question.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do not use an erasable pen or correction fluid.
- Do not write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 60.
- The number of marks for each question or part question is shown in brackets [].

This specimen paper has been updated for assessments from 2020. The specimen questions and mark schemes remain the same. The layout and wording of the front covers have been updated to reflect the new Cambridge International branding and to make instructions clearer for candidates.

For Examiner's Use	
Section A	
4	
5	
6	
Total	

This syllabus is regulated for use in England, Wales and Northern Ireland as a Cambridge International Level 3 Pre-U Certificate.

This document has 12 pages. Blank pages are indicated.

© UCLES 2018 [Turn over

Section A – Case Study

Read the passage carefully and answer all the questions.

You are advised to spend no more than 50 minutes on this section.

Type 2 diabetes – the growing threat

Diabetes mellitus currently affects at least 2.5 million people in the UK and is a condition in which the body is unable to maintain a normal blood glucose concentration. Many people who have no experience of diabetes think that the more common form is type 1, requiring insulin injections. Yet this is not the case. By far the more common is type 2, which represents approximately 85–90% of cases, and is on the increase. Originally thought of as affecting older people it is becoming increasingly common among the young. It is thought that obesity is an important risk factor. There is no entirely successful way of treating type 2 diabetes although it can be managed by control of diet, appropriate exercise and the use of medication.

Those with the condition, at least initially, produce insulin normally but certain body cells develop insulin resistance. This means that they do not respond to the hormone by taking up glucose from the blood rapidly enough to maintain a normal blood glucose concentration. The permeability of cell membranes is dependent on the presence of transporter protein molecules. Table 1.1 provides information about two types of such transporters, GLUT and SGLT. Table 1.1 distinguishes four types (isoforms) of GLUT.

Table 1.1

transporter group	type of mechanism	isoform	mainly present in	further information
		GLUT1	all cells	low-level basal glucose uptake required to sustain respiration
GLUT (glucose	facilitated diffusion	GLUT2	cells in small intestine lining, in the liver and in cells of kidney tubules	in the kidney tubule these transport glucose from cells lining the nephron into capillaries
transporters)		GLUT3	neurones	probably main glucose transporter in neurones
		GLUT4	adipose cells and striated muscle cells (skeletal and cardiac)	insulin-controlled glucose transporter
SGLT (sodium- glucose linked transporters)	secondary active transport along sodium gradient		cells lining the proximal tubule of nephrons	transport glucose directly from glomerular filtrate into cells lining nephron

Insulin is produced by the β cells of the islets of Langerhans within the pancreas. When the insulin concentration of the blood is low, GLUT4 molecules are removed from the cell membranes of adipose cells and skeletal muscle cells into vesicles in the cytoplasm. Except in the case of type 2 diabetes, an increase in blood insulin concentration means that insulin combines with specific sites on the cell surface membrane. This causes the GLUT4 molecules to be restored to the membrane, making it permeable to glucose. When blood insulin concentration falls, the GLUT4 molecules are removed from the membrane into cytoplasmic vesicles again.

On the onset of type 2 diabetes the patient's cells become insulin-resistant. Initially the pancreas responds by producing extra insulin. This only partially alleviates the problem of insulin resistance and, in time, overworking of the pancreatic β cells leads to their death and subsequently a reduction in insulin production. At this stage the patient may need to receive insulin injections, although this offers only a partial solution.

1	(a)	After a meal, blood glucose concentration rises above the target concentration $(4.5-5.5\mathrm{mmoldm^{-3}})$ at which it is normally maintained by homeostasis.
		With the help of Table 1.1, outline how the glucose concentration is reduced to normal in a person who does not have diabetes.
		[3]

(b) In the space below draw a simple, labelled diagram showing how protein transporter molecules may form part of a cell surface membrane.

(c)		plain how the uptake of glucose by cells in the proximal convoluted tubule differs from its ake by liver cells.
	•••••	[3]
		[3]
(d)	(i)	Suggest a mechanism by which the glucose transporter GLUT4 is restored to the membrane when insulin binds to the cell surface membrane.
		[1]
	(ii)	To what extent might the removal of GLUT4 from a muscle cell surface membrane render it impermeable to glucose?
		[1]
		[Total: 12]

BLANK PAGE

2 Table 2.1 presents the results of an experiment comparing rates of glucose production by a group of people with type 2 diabetes and a control group without the condition, during 23 hours of fasting.

Table 2.1

	rate of glucose product	tion per unit body mass g ⁻¹ min ⁻¹	oignificance level	
	patients with type 2 diabetes	control group	significance level	
total glucose production	11.1 ± 0.6	8.9 ± 0.5	p < 0.05	
glucose from hydrolysis of glycogen in the liver	1.3 ± 0.2	2.8 ± 0.7	p < 0.05	
glucose from gluconeogenesis	9.8 ± 0.7	6.1 ± 0.5	p < 0.01	

(a)	Discuss the conclusions which can be drawn from the data in Table 2.1.
	ro
	[6]

(b)	In individuals without diabetes, the blood glucose concentration in the renal vein is only slightly lower than in the renal artery.
	Explain why one might expect the glucose concentration of the blood in the renal vein to be much lower than in the renal artery and suggest why, in fact, the concentrations are almost identical.
	[4]
	[Total: 10]

3

It is possible to restore insulin secretion in a diabetic patient by transplanting a pancreas or isolated islets of Langerhans, but limited donor organs and risks involved restrict these therapies to a small

proportion of people with diabetes. Recent experiments suggest that it may, in future, be possible to treat diabetes with adult stem cells from the patient's own bone marrow.
Explain why such an approach, once perfected, is more likely to offer, at least initially, a treatmen for type 1 rather than type 2 diabetes and discuss why this approach may be preferable to the use of transplants or embryonic stem cells.

Section B - Synoptic Essay

Answer one question on the lined paper that follows.

You are advised to spend no more than 50 minutes on this section.

Choose **one** question from question 4, question 5 or question 6. Your answer should draw from a wide range of syllabus material and also demonstrate evidence of reading around the subject.

EITHER

4	'There is no evolutionary advantage in being multicellular'.	
	Discuss this view.	[30]
OR		
5	All living organisms need to synthesise ATP. Explain the similarities and differences borganisms in the ways in which this is achieved.	etween [30]
OR		
6	Why do people get heart disease and what should be done about it?	[30]

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of the University of Cambridge Local Examinations Syndicate (UCLES), which itself is a department of the University of Cambridge.