



Cambridge IGCSE™

CANDIDATE
NAME

CENTRE
NUMBER

--	--	--	--	--

CANDIDATE
NUMBER

--	--	--	--

PHYSICS

0625/62

Paper 6 Alternative to Practical

February/March 2021

1 hour

You must answer on the question paper.

No additional materials are needed.


INSTRUCTIONS

- Answer **all** questions.
- Use a black or dark blue pen. You may use an HB pencil for any diagrams or graphs.
- Write your name, centre number and candidate number in the boxes at the top of the page.
- Write your answer to each question in the space provided.
- Do **not** use an erasable pen or correction fluid.
- Do **not** write on any bar codes.
- You may use a calculator.
- You should show all your working and use appropriate units.

INFORMATION

- The total mark for this paper is 40.
- The number of marks for **each** question or part question is shown in brackets [].

This document has **16** pages. Any blank pages are indicated.



1 A student investigates the motion of an oscillating metre rule.

He uses the apparatus shown in Fig. 1.1.




Fig. 1.1

(a) The student ensures that the metre rule is horizontal.

Briefly describe how to check that the metre rule is horizontal. You may draw a diagram or draw on Fig. 1.1 if it helps to explain your answer.

- Measure the height of the metre rule from the horizontal bench from two different places. If equal, then ruler is horizontal. [1]

- (b) The student moves the stands so that the vertical threads are at the marks on the metre rule shown in Fig. 1.2.




Fig. 1.2

Calculate the distance d between the threads.

$$\begin{array}{r} -98 \\ \quad 2 \\ \hline 96 \end{array}$$

$d = \dots\dots\dots 96 \dots\dots\dots \text{cm}$ [1]

- (c) He twists the metre rule a small amount, as shown in Fig. 1.3, and then lets it go so that it oscillates in a rotating motion.




Fig. 1.3

He measures the time t for 5 complete oscillations of the metre rule.

$t = \dots\dots\dots 3.63 \dots\dots\dots \text{s}$

Suggest why it is useful to take a trial reading for this experiment.

- To check whether t is measurable for a given number of oscillations. [1]

- (d) The student carries out the same procedure for d values of 20.0 cm, 30.0 cm, 40.0 cm, 50.0 cm and 60.0 cm. His readings are shown in Table 1.1.

Table 1.1

d/cm	t/s	$\frac{1}{T}/\text{s}$
20.0	17.85	0.28
30.0	11.36	0.44
40.0	8.77	0.57
50.0	6.93	0.72
60.0	5.68	0.88

$$\frac{17.85}{5} = 3.57$$


$$\frac{1}{T} = \frac{1}{3.57} = 0.28$$

For distance $d = 20.0$ cm, calculate and record in Table 1.1, the value of $\frac{1}{T}$ where T is the time for 1 oscillation of the metre rule.

Use the value of time t from Table 1.1 and the equation $\frac{1}{T} = \frac{5}{t}$.

[1]

- (e) Plot a graph of distance d/cm (y-axis) against $\frac{1}{T}/\text{s}$ (x-axis).



[4]

- (f) Determine the gradient G of the graph. Show clearly on the graph how you obtained the necessary information.

$$G = \frac{\Delta y}{\Delta x} = \frac{57 - 21}{0.88 - 0.30} = 62$$

$$\frac{\text{cm}}{\frac{1}{\text{s}}} = \text{cm/s}$$

$$G = \dots\dots\dots 62 \text{ cm/s} \dots\dots\dots [1]$$

- (g) (i) Explain why it is more accurate to measure the time for 5 oscillations rather than for 1 oscillation.


- For 5 oscillations, the timing errors due to reaction time will have less effect compared for timing 1 oscillation. [1]

- (ii) Describe how the experiment could be improved to make the readings more reliable.

- Repeat each reading and calculate average value.

[1]

[Total: 11]



- 2 Students investigate the cooling of hot water in two different cups.

They use the apparatus shown in Fig. 2.1.

Cup A is made from thin plastic. The top of cup A has an inside diameter of 7 cm. Cup B is made from expanded polystyrene. The top of cup B has an inside diameter of 8 cm.




Fig. 2.1

- (a) (i) Record the room temperature θ_R shown on the thermometer in Fig. 2.1.

$$\theta_R = \dots\dots\dots 23^\circ\text{C} \dots\dots\dots [1]$$

- (ii) Describe **one** precaution that you would take to ensure that temperature readings in the experiment are as accurate as possible.

Wait until the mercury stops rising before taking reading. Also stir before reading, so that the heat is evenly (uniformly) distributed. [1]

- (b) A volume of 100cm^3 of hot water is poured into each cup and the initial temperature θ is recorded in Table 2.1. The temperature of the water in each cup is recorded every 30s. The values are shown in Table 2.1.

Table 2.1

	cup A	cup B
t/s	$\theta/^\circ\text{C}$	$\theta/^\circ\text{C}$
0	87.5	88.0
30	84.5	86.0
60	82.0	84.5
90	80.5	83.0
120	79.0	82.0
150	78.0	81.0
180	77.0	80.5

$$\begin{array}{r} 87.5 \\ - 88.0 \\ \hline 0.5^\circ\text{C} \end{array}$$

$$\begin{array}{r} 87.5 \\ - 77.0 \\ \hline 10.5^\circ\text{C} \end{array}$$

$$\begin{array}{r} 88.0 \\ - 80.5 \\ \hline 8.5^\circ\text{C} \end{array}$$

Complete the headings in Table 2.1.

[1]

- (c) Write a conclusion stating which cup, A or B, is the more effective in reducing the cooling rate of the hot water in this experiment.

Justify your answer by reference to the results.

- Cup B has a less cooling effect
 - Cup B had a temp decrease of 8.5°C
 compared to Cup A temp decrease of 10.5°C
 for the same time of 180s.

[2]

- (d) (i) Calculate x_A , the average cooling rate for cup A over the whole experiment. Use the readings for cup A from Table 2.1 and the equation

$$x_A = \frac{\theta_0 - \theta_{180}}{T}$$

where $T = 180\text{s}$ and θ_0 and θ_{180} are the temperatures at time $t = 0$ and at time $t = 180\text{s}$. Include the unit for the cooling rate.

$$\begin{aligned} x_A &= \frac{87.5 - 77}{180} \\ &= 0.058^\circ\text{C/s} \end{aligned}$$

$$x_A = \dots\dots\dots 0.058^\circ\text{C/s} \quad [2]$$

- (ii) Suggest an additional experiment to show how the lid affects the cooling rate of cup A.

Explain how to use the additional results to show the effect.

additional experiment *Repeat Cup A experiment
without a lid*

explanation *Calculate the cooling rate
and Subtract X_A .*

[2]


- (e) A student wishes to compare the effect of the materials of the cups on cooling rates. Suggest **two** variables that she should control to make this test fair.

1. *Same initial temperature*

2. *Same volume of water.*

[2]

[Total: 11]



- 3 A student investigates a resistor and a lamp connected in series. She uses the circuit shown in Fig. 3.1.




Fig. 3.1

- (a) The student moves the crocodile clip on the resistance wire so that the value of the potential difference V_L across the lamp is 2.0 V.

She measures the current I for the lamp and resistor in series.

She then connects the voltmeter to measure the potential difference V_R across the resistor.




Fig. 3.2




Fig. 3.3

Read, and record in Table 3.1, the values of I and V_R shown on the meters in Fig. 3.2 and Fig. 3.3.

[2]

- (b) The student repeats the steps in (a) for values of $V_L = 1.0\text{V}$ and $V_L = 0.5\text{V}$. Her readings are shown in Table 3.1.

Table 3.1

V_L/V	I/A	V_R/V	R_L/Ω	R_R/Ω
2.0	0.18	3.7	11.1	20.6
1.0	0.15	3.0	6.7	20.0
0.5	0.12	2.4	4.2	20.0

Calculate, and record in Table 3.1, the resistance of the lamp R_L for each value of V_L .

Use the values of V_L and I from Table 3.1 and the equation $R_L = \frac{V_L}{I}$.

Calculate, and record in Table 3.1, the resistance of the resistor R_R for each value of V_L .

Use the values of V_R and I from Table 3.1 and the equation $R_R = \frac{V_R}{I}$.

[2]

- (c) (i) Describe the pattern of any change in the value of R_L as V_L decreases.

- R_L decreases as V_L decreases.

[1]

- (ii) A student suggests that R_R should be constant.

State whether your results support this suggestion.

Justify your statement by reference to values from Table 3.1.

statement

Yes R_R is a constant

justification

the values of R_R are the same within limits of experimental accuracy.

[2]

- (d) A student wishes to determine the resistance of the lamp R_L when the potential difference across the lamp $V_L = 0.0\text{V}$.

Describe how the experiment can be extended to do this with the help of a suitable graph.

- Obtain more values of both V_L and R_L .
- Plot a R_L vs V_L graph.
- The graph cuts R_L axis at $V_L = 0.0$,
- So read R_L value where graph cuts axis. [2]

- (e) It is possible to use a variable resistor instead of a resistance wire to change the potential difference across the lamp.

Complete the circuit in Fig. 3.4 to show:

- a variable resistor used for this purpose
- the voltmeter connected to measure the potential difference across the resistor




Fig. 3.4

[2]

[Total: 11]

- 4 A student investigates the motion of a ball rolling down a slope.

Plan an experiment which enables him to investigate how **one** factor affects the average speed of the ball.

Average speed can be calculated using the equation:

$$\text{average speed} = \frac{\text{distance travelled}}{\text{time taken}}$$


The apparatus available includes:

balls of various sizes and materials
 a board which can act as a slope
 blocks to support one end of the board.

In your plan, you should:

- state a factor which can be measured
- list any additional apparatus needed
- explain briefly how to carry out the experiment including exactly which measurements are to be taken
- state the key variables to be kept constant
- draw a table, or tables, with column headings, to show how to display the readings (you are **not** required to enter any readings in the table)
- explain how to use the readings to reach a conclusion.

You may draw a diagram if it helps to explain your plan.



- The angle between the slope and the bench will be varied and measured.
- The mass of ball is same.
- The length of the slope is same
- Method:

Using a stop watch, measure time the ball takes at different angles to move down the slope.

- Measure the length of slope using a tape measure.
- Repeat each time and take average.
- Change the angle and repeat to measure time for the ball motion.
- Table

Angle/ $^{\circ}$	time/s

- Compare time readings in the table to see if change in angle changes the speed of the ball.

[7]

[Total: 7]