

0620 MCQ Answers

1-A	11-C	21-C	31-B
2-D	12-B	22-D	32-B
3-D	13-D	23-B	33-C
4-B	14-C	24-B	34-B
5-C	15-C	25-B	35-B
6-B	16-D	26-D	
7-	17-C	27-D	
8-	18-C	28-D	
9-C	19-D	29-D	
10-D	20-B	30-D	

0620 Theory Answers

Question 1

(c) $0.104/0.026$ [1]
 $n = 4$

Question 2

(c) mass of hydrated magnesium sulfate = 1.476 g
mass of barium sulfate formed = 1.398 g
the mass of one mole of BaSO_4 = 233 g
the number of moles of BaSO_4 formed = 0.006 [1]
the number of moles of $\text{MgSO}_4 \cdot x\text{H}_2\text{O}$ used in experiment = 0.006 [1]
the mass of one mole of $\text{MgSO}_4 \cdot x\text{H}_2\text{O}$ = $1.476/0.006 = 246$ g [1]
the mass of $x\text{H}_2\text{O}$ in one mole of $\text{MgSO}_4 \cdot x\text{H}_2\text{O}$ = $246 - 120 = 126$ g [1]
 $x = 126/18 = 7$ [1]
if x given without method = max 1
note: apply ecf but x must be an integer and less than 10

Question 3

(c) calculation:
Mr for NaHCO_3 = 84 g; Mr for Na_2O = 62 g; Mr for NaOH = 40 g
Mr for Na_2CO_3 = 106 g
(i) number of moles of NaHCO_3 used = $3.36/84 = 0.04$ [1]
(ii) if residue is Na_2O , number of moles of Na_2O = $2.12/62 = 0.034 / 0.03$
if residue is NaOH , number of moles of NaOH = $2.12/40 = 0.053 / 0.05$
if residue is Na_2CO_3 , number of moles of Na_2CO_3 = $2.12/106 = 0.02$ all three correct [2]
note: two correct = 1
(iii) equation 3 [1]
mole ratio 2:1 agrees with equation [1]

Question 4

(b) number of moles of HCl used = $0.04 \times 2 = 0.08$
number of moles CoCl_2 formed = 0.04

number of moles $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ formed = 0.04
mass of one mole of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ = 238 g
maximum yield of $\text{CoCl}_2 \cdot 6\text{H}_2\text{O}$ = 9.52g [4]
accept 9.5 g
mark ecf to moles of HCl
do not mark ecf to integers
to show that cobalt(II) carbonate is in excess
number of moles of HCl used = 0.08 must use value above ecf
mass of one mole of CoCO_3 = 119g
number of moles of CoCO_3 in 6.0g of cobalt(II) carbonate = $6.0/119 = 0.050$ [1]
reason why cobalt(II) carbonate is in excess $0.05 > 0.08/2$ [1]

Question 5

(d) (i) how many moles of H_2SO_4 were added = $0.02 \times 0.3 = 0.006$ [1]
(ii) how many moles of NaOH were used = $0.04 \times 0.2 = 0.008$ [1]
(iii) sulfuric acid [1]
only mark ecf if in accord with 1:2 ratio and with values from (i) and (ii).
reason $0.006 > 0.008/2$ [1]
for ecf mark candidate must use 1:2 ratio in answer
(iv) less than 7 [1]

Question 6

(b) (i) 80 cm³ of oxygen therefore 40 cm³ of methane [1]
 $40/60 \times 100 = 66.7\%$ [1]
accept 66 % and 67 %
no ecf
(ii) add sodium hydroxide(aq) / alkali [1]
carbon dioxide dissolves, leaving methane [1]

Question 7

(b) (i) add up to 5.8 g [1]
(ii) moles of C atoms = $2.4/12 = 0.2$
moles of H atoms = $0.2/1 = 0.2$
moles of O atoms = $3.2/16 = 0.2$
all three correct = 2 [2]
two correct = 1
empirical formula CHO [1]
(iii) $116/29 = 4$ [1]
 $\text{C}_4\text{H}_4\text{O}_4$ [1]
correct formula with no working scores both marks.
(iv) $\text{HOOCCH}=\text{CHCOOH}$ / $\text{CH}_2=\text{C}(\text{COOH})_2$ [2]

Question 8

(c) number of moles of FeSO_4 used = $9.12/152 = 0.06$ [1]
number of moles of Fe_2O_3 formed = 0.03^* [1]
mass of one mole of Fe_2O_3 = 160 g [1]
mass of iron(III) oxide formed = $0.03 \times 160 = 4.8$ g [1]
number of moles of SO_3 formed = 0.03 [1]
volume of sulfur trioxide formed = $0.03 \times 24 = 0.72$ dm³ [1]
If mass of iron(III) oxide greater than 9.12 g, then only marks 1 and 2 available
Apply ecf to number of moles of Fe_2O_3^* when calculating volume of sulfur trioxide.

Do not apply ecf to integers

Question 9

7 (a) (i) 35 cm³ [1]
40 cm³ [1]

Question 10

(b) (i) 7.7% [1]
(ii) for any number: equal number ratio [2]
for example 1:1 or 6:6
(iii) empirical formula is CH [1]
molecular formula is C₆H₆ [1]
no e.c.f., award of marks not dependent on (ii)

Question 11

(c) (i) 196 [1]
(ii) $112/196 \times 100$ [1]
= 57(.1)% ACCEPT 57 to nearest whole number [1]
mark e.c.f. to (c)(i) provided percentage not greater than 100%
ONLY ACCEPT 112/answer (c)(i) $\times 100$
otherwise [0]

Question 12

(ii) mass of one mole of CaCO₃ = 100
number of moles of CaCO₃ = $0.3/100 = 0.003$ [1]
moles of HCl = $5/1000 \times 1 = 0.005$ [1]
reagent in excess is CaCO₃ [1]
ecf from above
would need 0.006 moles of HCl
or hydrochloric acid only reacts with 0.0025 moles of CaCO₃ [1]
NOTE this mark needs to show recognition of the 1:2 ratio
(iii) mark ecf to (ii), that is from moles of limiting reagent in (ii)
moles of CO₂ = $0.005 \times 0.5 \times 24 = 0.06 \text{ dm}^3$ [1]
NOT cm³ unless numerically correct. 60 cm³
Ignore other units
NOTE If both number of moles integers then no ecf for (ii) and (iii)

Question 13

(a)
copper iron sulphur
composition by
mass/g
(4.80) (4.20) 4.8 [1]
number of moles
of atoms
0.075 0.075 0.15 [1]
simplest mole ratio
of atoms
1 1 2 [1]
[3]
The empirical formula is CuFeS₂ [1]

Question 14

(b) (i) 100 [1]
56 ignore units in both cases [1]
(ii) 7.00kg is 1/8 of 56 [1]
1/8 of 100kg is 12.5kg [1]
Give both marks for correct answer without explanation.
Ignore missing units
but penalise wrong units

Question 15

Question 6
(a)(i) moles of NiCO₃ reacted = 0.08 [1]
mass of nickel carbonate reacted = 9.52 g [1]
mass of nickel carbonate unreacted = 2.48 g [1]
(ii) maximum number of moles of hydrated salt = 0.08 [1]
maximum mass of salt = $0.08 \times 281 = 22.48 \text{ g}$ [1]
percentage yield $10.4/22.48 \times 100 = 46.3\%$ [1]

Question 16

Mark consequentially to any error but not involving simple integers
There has to be some evidence that the candidate has attempted to work through the calculation and not merely inserted whole numbers.
For example 2, 1, 160 or 1, 0.5, 80
number of moles of Fe₂(SO₄)₃ = 1/40 or 0.025
number of moles of Fe₂O₃ formed = 1/40 or 0.025
mass of iron(III) oxide formed = $0.025 \times 160 = 4\text{g}$
number of moles of SO₃ produced = 3/40 or 0.075
volume of sulphur trioxide at r.t.p. = $0.075 \times 25 = 1.8\text{dm}^3$ [5]

Question 17

(d) the number of moles of SO₂ in the mixture = 0.125
the number of moles of Cb in the mixture = 0.2
cond reagent was not in excess? SO₂
cond moles of SO₂Cb formed = 0.125
cond the mass of sulphuryl chloride formed = 16.9g [5]

Question 18

(f) (i) $11.5/23 = 0.5$ [1]
(ii) 0.25 [1]
conseq to (i)
...
(iii) $0.25 \times 32 = 8 \text{ g}$ [1]
conseq
(iv) 2.0 g [1]
only conseq to (iii) if answer to (iii) is less than 10
NB If (ii) is 0.3(125), no excess is possible, (iv) ZERO

Question 19

(c) (i) copper sulphate or anhydrous copper sulphate [1]
accept "unhydrated"
NOT formula
(ii) goes blue or becomes hot or steam [1]

(iii) copper oxide [1]
(iv) $5/250 = 0.02$ moles
Mr=80
 $80 \times 0.02 = 1.6$ g
NB (iv) to be marked conseq to (iii)
Correct answer no working ONLY [1]

Question 20

(e) (i) percentage of oxygen = 31.6 % [1]
(ii) calculate the number of moles of atoms for each element
number of moles of Ti = $31.6/48 = 0.66$
number of moles of O = $31.6/16 = 1.98$ accept 2 [1]
both correct for one mark
(iii) the simplest whole number ratio for moles of atoms:
Fe : Ti : O
1 1 3 [1]
(iv) formula is FeTiO₃ accept TiFeO₃ [1]
must be whole numbers from (iii) or cancelled numbers from
(iii)
mark ecf throughout

Question 21

(ii) Volume ratio
C_x
Hy(g) + O₂(g) → CO₂(g) + H₂O(l)
20 160 100 all in cm³
1 8 5 mole ratio
C₅
H₁₂ + 8O₂ → 5CO₂ + 6H₂O
For evidence of method (1)
for equation as above (2) [3]

Question 22

(c) (i) (to prove) all water driven off or evaporated or boiled /
no water remains / to
make salt anhydrous (1)
(ii) $m_1 - m_2 = \text{mass of water}$ (1)
(calculate) moles of water AND moles of hydrated or
anhydrous salt (1)
1:1 ratio / should be equal (1) [3]

Question 23

(d) number of moles of O₂ formed = $0.096 / 24 = 0.004$ (1)
number of moles of H₂O₂ in 40 cm³ of solution = $0.004 \times 2 =$
0.008 (1)
concentration of the hydrogen peroxide in mol / dm³ = 0.008
/ $0.04 = 0.2$ (1) [3]

Question 24

8 (a) (i) (the number of particles which is equal to the number
of atoms in) 12 g of carbon 12
or
the mass in grams which contains the Avogadro's constant
number of particles
or
Avogadro's constant or 6 to 6.023×10^{23} of atoms / ions /
molecules / electrons /

particles
or
(the amount of substance which has a mass equal to) its
relative formula mass / relative
atomic mass / relative molecular mass in grams
or
(the amount of substance which has a volume equal to) 24
dm³ of a gas at RTP
[1]
(ii) (Avogadro's constant is the) number of particles / atoms /
ions / molecules in one mole of
a substance
or
the number of carbon atoms in 12 g of C(12).
or
the number of particles / molecules in 24 dm³ of a gas at RTP
or
6 to 6.023×10^{23} (particles / atoms / ions / molecules /
electrons) [1]
(b) CH₄ and SO₂ [1]
 $2/16 = 1/8$ or 0.125 moles of CH₄ AND $8/64 = 1/8$ or 0.125
moles of SO₂
(c) (i) $4.8/40 = 0.12$ moles of Ca
 $3.6/18 = 0.2$ moles of H₂O both correct [1]
(ii) Ca is in excess (no mark) (because 0.12 moles of Ca need)
0.24 moles / 4.32 g of H₂O
to react [1]
there is not enough / there are 0.2 moles / 3.6 g of H₂O [1]
or
Ca is in excess (no mark) (because 0.2 moles / 3.6 g of water
will react with)
0.1 moles / 4.0 g of Ca [1]
there is more than that / there are 0.12 moles / 4.8 g of Ca [1]
or
Ca is in excess (no mark) because the mole ratio Ca:H₂O is 3:5
/ mass ratio 4:3 [1]
which is bigger than the required mole ratio of 1:2 / mass ratio
10:9 [1]
or
Ca is in excess (no mark) because the mole ratio H₂O:Ca is 5:3
/ mass ratio 3:4 [1]
which is smaller than the required mole ratio of 2:1 / mass
ratio 9:10 [1]
(iii) $0.02 \times 40 = 0.8$ (g) [1]

Question 25

(d) volume of oxygen used = 150 cm³
volume of carbon dioxide formed = 100 cm³ [1]
any equation of the combustion of an alkene
e.g. $2C_5H_{10} + 15O_2 \rightarrow 10CO_2 + 10H_2O$
formulae [1]
COND balancing

Question 26

(b) number of moles of HCl = $0.020 \times 2.20 = 0.044$ [1]
number of moles of LiOH = 0.044
concentration of LiOH = $0.044/0.025 = 1.769$ (mol / dm³) [1]

accept 1.75 to 1.77 need 2 dp
correct answer scores = 2
(c) (for $\text{LiCl}\cdot 2\text{H}_2\text{O}$)
mass of one mole = 78.5 [1]
percentage water = $36 / 78.5 \times 100$ [1]
45.9 so is $\text{LiCl}\cdot 2\text{H}_2\text{O}$ [1]
only award the marks if you can follow the reasoning and it gives 45.9% of water
note: if correct option given mark this and ignore the rest of the response
allow: max 2 for applying a correct method to another hydrate, [1] for the method and [1] for the correct value, working essential

Question 27

(e) if C_5H_{10} is given award 3 marks;;; [3]
if $\text{C}_{10}\text{H}_{20}$ is given award 2 marks;;
if 1:7.5:5 / 2:15:10 is given award 2 marks;;
in all other cases a mark can be awarded for moles of O_2 (= $2.4/32$ =) 0.075 AND moles of CO_2 (= $2.2/44$ =) 0.05;
 $2\text{C}_5\text{H}_{10} + 15\text{O}_2 \rightarrow 10\text{CO}_2 + 10\text{H}_2\text{O}$ [1]
accept: multiples including fractions
allow: ecf for correct equation from any incorrect alkene

Question 28

(b) moles of Fe = $51.85/56 = 0.926$ (0.93); [1]
moles of O = $22.22/16 = 1.389$ (1.39); [1]
moles of $\text{H}_2\text{O} = 16.67/18 = 0.926$ (0.93); [1]
if given as 0.9 1.4 0.9
three of the above correct = [2]
two of the above correct = [1]
simplest whole number mole ratio Fe : O : H_2O is 2 : 3 : 2 / $\text{Fe}_2\text{O}_3\cdot 2\text{H}_2\text{O}$; [1]
allow: ecf for a formula based on an incorrect whole number ratio

Question 29

8 (a) (i) (to avoid) carbon monoxide formation/so complete combustion occurs/avoid incomplete combustion So that CO_2 is produced [1]
CO does not dissolve/react with alkali [1]
(ii) CO_2 is acidic [1]
(iii) volume of gaseous hydrocarbon 20 cm^3
volume of oxygen used = 90 cm^3 [1]
volume of carbon dioxide formed = 60 cm^3 [1]
no mark for 20 cm^3 of hydrocarbon.
(iv) $2\text{C}_3\text{H}_6(\text{g}) + 9\text{O}_2(\text{g}) \rightarrow 6\text{CO}_2(\text{g}) + 6\text{H}_2\text{O}(\text{l})$ [1]
OR ... $\text{C}_3\text{H}_6(\text{g}) + 9/2\text{O}_2(\text{g}) \rightarrow 3\text{CO}_2(\text{g}) + 3\text{H}_2\text{O}(\text{l})$
 C_3H_6 [1]
 C_3H_6 can be given in the equation for the second mark

Question 30

7 (a) metal A is magnesium [1]
cond most reactive or fastest reaction [1]
metal B is aluminium [1]

cond faster reaction after removal of oxide layer / it would give more hydrogen / aluminium more reactive than zinc [1]
metal C is zinc [1]
zinc least reactive [1]
NOTE MAX [5]
If you encounter different reasoning which is correct, please award the appropriate marks.
(b) for magnesium and zinc same volume of hydrogen [1]
because both have valency of 2 / 1 mole of metal gives 1 mole of hydrogen / 1 mole of metal reacts with 2 moles of acid [1]
bigger volume for aluminium because its valency is 3 / 1 mole of metal gives 1.5 moles of hydrogen / 1 mole of metal reacts with 3 moles of acid [1]
If you encounter different reasoning which is correct, please award the appropriate marks.
accept balanced equations
accept ionic charges as alternative to valency

Question 31

(d) (i) the reaction is exothermic / reaction produces heat/energy [1]
all the sodium hydroxide used up/neutralised / reaction has stopped [1]
(ii) adding colder acid / no more heat produced [1]
if not given in (d)(i) any comments such as "reaction has stopped" can gain mark
(iii) 1.33 / 1.3 / 1.3333 (mol/ dm^3) scores both marks [2]
not 1.34
for a correct method – $M_1 V_1$ / moles of $\text{NaOH} = 0.02$
with an incorrect answer only [1]

Question 32

(c) if the final answer is between 86–89% award all 4
if the final answer is between 66–67% award 3 marks (Mr of 32 must have been used)
for all other answers marks can be awarded using the mark scheme as below and applying ecf if necessary
number of moles of O_2 formed = $0.16/24 = 0.0067/0.00667$ or $1/150$
number of moles of $\text{Pb}(\text{NO}_3)_2$ in the sample = $0.0133/0.013$ or $1/75$
mass of one mole of $\text{Pb}(\text{NO}_3)_2 = 331$ g
mass of lead(II) nitrate in the sample = 4.4(1) g
percentage of lead(II) nitrate in sample = 88.3% (allow 88–89) [4]
mark ecf in this question but not to simple integers
if mass of lead(II) nitrate > 5.00 only marks 1 and 2 available
If divides by 32 (not 24) only last 3 marks can score consequentially

Question 33

(a) $72/24 = 3$ and $28/14 = 2$ [1]
 Mg_3N_2 [1]
accept just formula for [2] even with incorrect or no working

NOT ecf

(b) $Al_4C_3 + 12H_2O = 4Al(OH)_3 + 3CH_4$ [2]

For Al_4C_3 ONLY [1]

(c) (i) silicon is limiting reagent [1]

0.07 moles of Si and $25/160 = 0.156$ moles of Br_2 [1]

because $0.14 (2 \times 0.07) < 0.156$ [1]

If 80 used to find moles of Br_2 the mark 1 and 3 still available arguments based on masses can be used

(ii) 0.07 [1]

NOT ecf

Question 34

(b) number of moles of NaOH used = $0.025 \times 2.24 = 0.056$ [1]

maximum number of moles of $Na_2SO_4 \cdot 10H_2O$ that could be formed = 0.028 [1]

mass of one mole of $Na_2SO_4 \cdot 10H_2O = 322g$

maximum yield of sodium sulphate – 10 - water = 9.02g [1]

percentage yield = 42.8% [1]

mark ecf but NOT to simple integers

if ecf marking, mark to at least one place of decimals

if percentage > 100% then 3/4 maximum

Question 35

(d) 100g of fat react with 86.2g of iodine

884g of fat react with 762 g of iodine [1]

limit 762 x 2

one mole of fat reacts with 762/254 moles of iodine molecules

one mole of fat reacts with 3 moles of iodine molecules [1]

number of double bonds in one molecule of fat is 3 [1]

limit 6

consequential marking allowed provided the number of double bonds is an integer.

Question 36

(d) moles of $CH_3-CH=CH_2$ reacted = $1.4/42 = 0.033$ [1]

conseq

maximum moles of $CH_3-CH(I)-CH_3$ that could be formed = 0.033 [1]

conseq

maximum mass of 2-iodopropane that could be formed = 5.61 g [1]

accept $170 \times 0.033 = 5.61$ and $170 \times 0.033333 = 5.67$

conseq unless greater than 100%

percentage yield $4.0/5.67 \times 100 = 70.5\%$ [1]

Do not mark consequently to a series of small integers. There has to be

a serious attempt to answer the question, then consequential marking is appropriate.

Question 37

(d) mass of one mole of $CaSO_4 = 136$

moles of $CaSO_4$ in 79.1g = 0.58 accept 0.6 [1]

moles of H_2O in 20.9 g = 1.16 accept 1.2 [1]

conseq $x = 2 \times$ given as an integer [1]

Question 38

(c) $I_2 + 3Cl_2 = 2ICl_3$ [2]

For having either reactants or products correct ONLY [1]

Question 39

skip

Question 40

(c) (i) number of moles $CO_2 = 0.24/24 = 0.01$

conseq number of moles of $CaCO_3$ and $MgCO_3 = 0.01$

conseq number of moles of $CaCO_3 = 0.005$ [3]

(ii) Calculate the volume of hydrochloric acid, 1.0 mole/dm³, needed to react with one tablet.

number of moles of $CaCO_3$ and $MgCO_3$ in one tablet = 0.01

Expect same as answer to (c)(i). NO marks to be awarded. Just mark

consequentially to this response

conseq number of moles of HCl needed

to react with one tablet = 0.02

conseq volume of hydrochloric acid, 1.0 mole/dm³, needed to react with one

tablet = 0.02 dm³ or 20 cm³

[1]

[1]

Question 41

(c) number of moles of HCl in 50 cm³ of acid, concentration

2.2 mol/dm³ = 0.11 [1]

maximum number of moles of $CoCl_2 \cdot 6H_2O$ which could be formed = 0.055 [1]

mass of 1 mole of $CoCl_2 \cdot 6H_2O = 238 g$

maximum yield of $CoCl_2 \cdot 6H_2O = 13.09 g$ [1]

percentage yield = 48.2% or ecf mass of $CoCl_2 \cdot 6H_2O$

above/ $13.09 \times 100\%$ to 1

dp [1]

Question 42

(b) (i) 14.3 [1]

(ii) $85.7 \div 12$ and $14.3 \div 1$ or 7.14 and 14.3 [1]

ratio 1:2 [1]

CH_2 [1]

note: Award all 3 marks for correct answer

allow: alternative working e.g.

$85.7 \times 84 \div 100$ and $14.3 \times 84 \div 100$ or $71.988/72$ and

$12/12.012$ [1]

6:12 or ratio 1:2 [1]

CH_2 [1]

(iii) C_6H_{12} [1]

Question 43

(iii) $M_1 = 2.07$ Allow 2.1 or 2.0666...7

$M_2 = 62.8.g$

$M_3 = (M_2/152 =) 0.41(3)$

$M_4 (=M_1/M_3)$ rounded to the nearest whole number $\times = 5$ [4]

Question 44

(ii) number of moles of ethanoic acid = 0.1 [1]
number of moles of ethanol = 0.12(0) [1]
the limiting reagent is ethanoic acid [1]
number of moles of ethyl ethanoate formed = 0.1 [1]
maximum yield of ethyl ethanoate is 8.8 g [1]

(c) BrF₃ / F₃Br; [1]
BrF₅ / F₅Br; [1]

Question 45

(ii) mass of AgNO₃ needed is $170 \times 0.2 \times 0.1 = 3.4\text{g}$ [2]
NOTE: if answer given is 34 they have omitted 0.1
ALLOW: (1) ecf
(iii) number of moles of AgNO₃ used = $0.02 \times 0.2 = 0.004$ [1]
number of moles of Ag₂CrO₄ formed = 0.002 [1]
mass of one mole of Ag₂CrO₄ = 332g
mass of Ag₂CrO₄ formed = 0.664g [1]
NOTE: use ecf when appropriate

Question 46

(c) number of moles of CO₂ formed = $2.112 / 44 = 0.048$ [1]
number of moles of H₂O formed = $0.432 / 18 = 0.024$ [1]
x = 2 and y = 1 NOT: ecf from this line
formula is 2PbCO₃.Pb(OH)₂ / Pb(OH)₂. 2PbCO₃ [1]

Question 47

(d) number of moles of HCl in 40 cm³ of hydrochloric acid,
concentration 2.0 mol / dm³ = $0.04 \times 2.0 = 0.08$ [1]
maximum number of moles of CO₂ formed = 0.04 [1]
mass of one mole of CO₂ = 44 g [1]
maximum mass of CO₂ lost = $0.04 \times 44 = 1.76$ g [1]

Question 48

(b) (i) $(97.4 / 75 =)$ 1.3 and $(2.6 / 1 =)$ 2.6; [1]
empirical formula AsH₂; [1]
note: correct formula with no working = [1]
(ii) As₂H₄; [1]
(iii) H₂As–AsH₂ / AsH₂–AsH₂; [1]

Question 49

(d) number of moles of Na₂SO₃ = $3.15 / 126 = 0.025$ [1]
number of moles of SO₂ formed = 0.025 [1]
volume of SO₂ = $0.025 \times 24 = 0.6$ dm³/litres or 600 cm³ [1]
allow: ecf
for 1.6 g of SO₂ [1] only
If used 22.4 max [2]
note: need correct units for last mark

Question 50

(c) number of moles of HCl used = $0.05 \times 2 = 0.1$ [1]
number of moles of SrCl₂.6 H₂O which could be formed. =
0.05 [1]
mass of one mole of SrCl₂.6H₂O is 267 g
theoretical yield of SrCl₂.6H₂O = $0.05 \times 267 = 13.35$ g [1]
percentage yield = $6.4 / 13.35 \times 100 = 47.9\%$ [1]
accept: 48%
allow: ecf

Question 51